

THE ASYMPTOTICAL BEHAVIOR OF SPECTRAL FUNCTION OF ONE FAMILY OF DIFFERENTIAL OPERATORS.

ALEXANDER S. PECHENTSOV AND ANTON YU. POPOV
MOSCOW STATE UNIVERSITY, RUSSIA

In $L_2[0, +\infty)$ consider operator l , defined by differential expression:

$$ly(x) = -y''(x) + q(x)y(x)$$

and boundary condition

$$y(0) \cos \alpha + y'(0) \sin \alpha = 0.$$

Assume that $\alpha \in R$, q is a function continuous on $[0, +\infty)$, and having real values. Denote by $\varphi(x, \lambda)$ and $\theta(x, \lambda)$ the solutions to equation $ly = \lambda y$ with initial conditions

$$\varphi(0, \lambda) = \sin \alpha, \quad \varphi'(0, \lambda) = -\cos \alpha, \quad \theta(0, \lambda) = \cos \alpha, \quad \theta'(0, \lambda) = \sin \alpha.$$

Functions φ, θ and $\rho, m, \hat{f}$ (we will introduce them later) depend, naturally, on α , but we will omit the argument α , for not to overload the text with notations.

It is well known the Weyl theorem on representation of arbitrary function $f \in L_2[0, +\infty)$ as an integral along the spectrum of operator L .

Theorem [1-3]. *There exists such a non decreasing and bounded from below on $\mathbb{R}$ function ρ , for which the following statements take place:*

1. $\exists \hat{f}(\lambda) = \lim_{L_2(R, d\rho) \ n \rightarrow \infty} \int_0^n f(x) \varphi(x, \lambda) dx = \int_0^{+\infty} f(x) \varphi(x, \lambda) dx$,
and the function f is considered as a limit in $L_2[0, +\infty)$ of "Fourier transform by measure $d\rho$ " of $\hat{f}$:

$$f(x) = \lim_{L_2(R^+) \ n \rightarrow \infty} \int_{-n}^n \hat{f}(\lambda) \varphi(x, \lambda) d\rho(\lambda) = \int_{-\infty}^{+\infty} \hat{f}(\lambda) \varphi(x, \lambda) d\rho(\lambda).$$

2. If $\sin \alpha \neq 0$, then for function $m(z)$ defined in half-plane $\text{Im } z > 0$ by the equality

$$m(z) = -\text{ctg } \alpha + \int_{-\infty}^{+\infty} \frac{d\rho(\lambda)}{z - \lambda}, \quad (1)$$

we have:

$$\theta(x, z) + m(z)\varphi(x, z) \in L_2[0, +\infty) \quad \forall z : \text{Im } z > 0.$$

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$

3. For an arbitrary Borel set $E \subset R$, not including the spectrum of operator L , we have: $\int_E d\rho(\lambda) = 0$.

The statement 3 of Theorem shows, that the measure $d\rho$ is concentrated on the spectrum of operator L . The function $\rho(\lambda)$ is called the spectral function, and $m(z)$ is called the function of Weyl-Titchmarsh of operator L . It is easy to see from the representation (1) that the function $m(z)$ is analytical in the upper half-plane and that if on certain interval $(a, b) \subset R$ there exists

$$\lim_{y \rightarrow +0} m(\lambda + iy) = m(\lambda) \in C(a, b),$$

then $\rho(\lambda)$ has on this interval continuous derivative, which is connected to $m(\lambda)$ by the correspondence:

$$\rho'(\lambda) = \frac{-\operatorname{Im} m(\lambda)}{\pi}, \lambda \in (a, b). \quad (2)$$

The investigation of properties of function $p(\lambda)$, and its definition via the potential q and the number α , defining the operator L , is, as a rule, a very difficult problem. There are well known the general theorems of B.M.Levitan and V.A.Marchenko (see [4]—[6] on evaluation of $\rho(\lambda)$ as $\lambda \rightarrow \pm\infty$:

$$\rho(\lambda) - \rho(-\infty) = o(\exp(-a\sqrt{|\lambda|})), (\lambda \rightarrow -\infty) \forall a > 0, \quad (3)$$

$$\rho(\lambda) = \frac{2\sqrt{\lambda}}{\pi \sin^2 \alpha} + \rho(-\infty) + \frac{\cos \alpha}{\sin^2 \alpha} + o(1) (\lambda \rightarrow +\infty).$$

(it is assumed, that the norming $\rho(0) = 0$ is performed.)

The simplest case of $q(x) \equiv 0$ was studied by Titchmarsh in [7]. It turned out, that for $\lambda > 0$

$$\rho'(\lambda) = \frac{\sqrt{\lambda}}{\pi(\lambda \sin^2 \alpha + \cos^2 \alpha)},$$

and for $\lambda < 0$ both of the cases $\operatorname{ctg} \alpha > 0$ and $\operatorname{ctg} \alpha < 0$ are strongly different:

$$\rho'_\alpha(\lambda) \equiv 0, \lambda < 0, \operatorname{ctg} \alpha < 0,$$

$$\rho'_\alpha(\lambda) = \frac{2 \operatorname{ctg} \alpha}{\sin^2 \alpha} \delta(\lambda - \lambda_0), \lambda < 0, \operatorname{ctg} \alpha > 0, \quad (4)$$

where $\lambda_0 = -\operatorname{ctg}^2 \alpha$, δ is delta-function of Dirac.

In this work we study the case of $\operatorname{ctg} \alpha > 0$ and analyse the behavior of the derivative of spectral function on $(-\infty, 0)$ for potentials of special type.

The following problem of common form is interesting for our study:

Let $Q \in C(0, +\infty)$. Denominate as $\rho_\alpha(\varepsilon, Q, \lambda)$ the spectral function of the operator, defined by differential expression:

$$-y''(x) + \varepsilon Q(x)y(x)$$

and boundary condition

$$y(0) \cos \alpha + y'(0) \sin \alpha = 0, \operatorname{ctg} \alpha > 0.$$

Is it true that on $(-\infty, 0)$ the family of distributions $\rho'_\alpha(\varepsilon, Q, \lambda)$ converges as $\varepsilon \rightarrow 0$ (in any "reasonable" sence) to the derivative of spectral function (4) of non-perturbed operator $-y''$? The positive solution of this problem would allow us to write for functions $f \in L_2[0, +\infty)$ the "approximate spectral expansion" of the following kind:

$$f(x) = \int_{\lambda_0 - \beta}^{\lambda_0 + \beta} + \int_{-\beta}^{+\infty} \hat{f}(\lambda) \varphi_\alpha(x, \lambda) d\rho_\alpha(\varepsilon, Q, \lambda)$$

($\beta = \beta(\varepsilon) \downarrow 0$), omitting the integral on the complement to small neighborhood of the spectrum of non-perturbed operator. As we know, such investigations have not been performed before.

Titchmarsh in [7] have expressed through Gankel functions of the first type the Weyl-Titchmarsh function $m(\lambda, \varepsilon)$ of operator

$$-y'' - \varepsilon xy, \quad \varepsilon > 0, \quad y(0) \cos \alpha + y'(0) \sin \alpha = 0, \quad \text{ctg } \alpha > 0. \quad (5)$$

He proved, that for all $\lambda \in \mathbb{R}$

$$m(\lambda, \varepsilon) = \frac{H_{1/3}^{(1)}(A) \sin \alpha - \sqrt{\lambda} H_{-2/3}^{(1)}(A) \cos \alpha}{H_{1/3}^{(1)}(A) \cos \alpha + \sqrt{\lambda} H_{-2/3}^{(1)}(A) \sin \alpha}, \quad (6)$$

where $A = \frac{2\lambda^{3/2}}{3\varepsilon}, \lambda^{3/2} = -i|\lambda|^{3/2}$ with $\lambda < 0, H_p^{(1)}$ are Hankel functions of the first type.

Using the representation (6) it was proved in [6], that in certain vicinity of the point λ_0 the functions $m(\lambda, \varepsilon)$ have only one pole $\lambda_0(\varepsilon)$ with the asymptotics

$$\lambda_0(\varepsilon) = \lambda_0 - \frac{\varepsilon}{2} \text{tg } \alpha + O(\varepsilon^2), \quad (\varepsilon \rightarrow +0),$$

and its imaginary part satisfies the inequality:

$$-\exp\left(-\frac{\text{ctg}^3 \alpha}{\varepsilon}\right) \leq \text{Im } \lambda_0(\varepsilon) < 0.$$

But the behavior of $\rho'(\lambda, \varepsilon)$ as $\lambda \in (-\infty, 0)$ and $\varepsilon \rightarrow +0$ was not studied by Titchmarsh. We could solve this problem in certain sence. In spite of the existence of explicit formula for $\rho'(\lambda, \varepsilon)$ (taking into account (2) we have $\rho'(\lambda, \varepsilon) = -\frac{\text{Im } m(\lambda, \varepsilon)}{\pi}$), due to the fact that from (6) it follows, that the function $m(\lambda, \varepsilon)$ is analytical in closed upper half-plane except one point 0) this was not easily to realize. The matter is that in putting in (6) known asymptotic series for $H_{1/3}^{(1)}(A)$ and $H_{-2/3}^{(1)}(A)$ as $A \rightarrow -i\infty$ (along negative part of imaginary axis), then for $m(\lambda, \varepsilon)$ we obtain the asymptotic series, all members of which are real. Therefore, mentioned series does not give information concerning $\rho'(\lambda, \varepsilon)$.

We have obtained the following representation for function $\rho'(\lambda, \varepsilon)$, which takes place for any $\lambda < 0$ and $\varepsilon > 0$:

$$\rho'(\lambda, \varepsilon) = \frac{\pi^{-1} \tau (\beta_{1,1}(a) \beta_{2,2}(a) + \beta_{1,2}(a) \beta_{2,1}(a))}{(\beta_{1,1}(a) \cos \alpha - \beta_{1,2}(a) \tau \sin \alpha)^2 + (\beta_{2,1}(a) \cos \alpha + \beta_{2,2}(a) \tau \sin \alpha)^2}, \quad (7)$$

where $\tau = \sqrt{-\lambda}$, $a = |A| = \frac{2\tau^2}{3\varepsilon}$,

$$\beta_{1,1}(a) = 1 + \frac{\sqrt{3}}{2} \frac{B_{1/3}^-(a)}{\Omega_{1/3}(a)}, \quad \beta_{1,2}(a) = \frac{\Omega_{2/3}(a)}{\Omega_{1/3}(a)} + \frac{\sqrt{3}}{2} \frac{B_{2/3}^-(a)}{\Omega_{1/3}(a)},$$

$$\beta_{2,1}(a) = \frac{B_{1/3}^+(a)}{2\Omega_{1/3}(a)}, \quad \beta_{2,2}(a) = \frac{B_{2/3}^+(a)}{2\Omega_{1/3}(a)},$$

$$\Omega_p(a) = \frac{1}{\pi} \int_0^\pi \exp(a \cos t) \cos(pt) dt,$$

$$B_p^-(a) = \frac{1}{\pi} \int_0^{+\infty} \exp(-a \operatorname{ch} t) \operatorname{sh}(pt) dt,$$

$$B_p^+(a) = \frac{1}{\pi} \int_0^{+\infty} \exp(-a \operatorname{ch} t) \operatorname{ch}(pt) dt.$$

The representation (7) allows us to prove the following theorems 1-3:

Theorem 1. *For any $\varepsilon > 0$ the following asymptotics takes place:*

$$\rho'(\lambda, \varepsilon) \sim \frac{\exp(-\frac{4}{3} \frac{|\lambda|^{3/2}}{\varepsilon})}{\pi \sqrt{-\lambda} \sin^2 \alpha}, \quad (\lambda \rightarrow -\infty).$$

Theorem 2. *There exist positive constants c_1 and c_2 , effectively dependent on α , such that with $\varepsilon \in (0, c_1)$ take place evaluations:*

$$\rho'(\lambda, \varepsilon) = O(|\lambda|^{-1/2} \exp(-\frac{4|\lambda|^{3/2}}{3\varepsilon})), \quad \lambda \in (-\infty, -2 \operatorname{ctg}^2 \alpha),$$

$$\rho'(\lambda, \varepsilon) = O(|\lambda|^{1/2} \exp(-\frac{4|\lambda|^{3/2}}{3\varepsilon})), \quad \lambda \in (-\frac{1}{2} \operatorname{ctg}^2 \alpha, -c_2 \varepsilon^{2/3}),$$

$$\rho'(\lambda, \varepsilon) = O(\varepsilon^{1/3}), \quad \lambda \in [-c_2 \varepsilon^{2/3}, 0).$$

Constants in symbols O depend only on α and are effective.

Denote $T(\lambda, \varepsilon) = \beta_{1,1}(a) \cos \alpha - \beta_{1,2}(a) \tau \sin \alpha$. The answer to the question on the behavior of $\rho'(\lambda, \varepsilon)$ on the "critical" segment $I = [-2 \operatorname{ctg}^2 \alpha, -\frac{1}{2} \operatorname{ctg}^2 \alpha]$ gives the following

Theorem 3. *There exists such a positive constant c_3 , depending effectively on α , that for any $\varepsilon \in (0, c_3)$ the following statements take place:*

1) *Function $T(\lambda, \varepsilon)$ has on segment I unique zero $\lambda_1(\varepsilon)$ with asymptotics:*

$$\lambda_1(\varepsilon) = \lambda_0 - \frac{\varepsilon}{2} \operatorname{tg} \alpha + O(\varepsilon^2).$$

2) *For any $d > 0, \lambda \in I \setminus I_{d,\varepsilon}$ ($I_{d,\varepsilon} = [\lambda_1(\varepsilon) - d, \lambda_1(\varepsilon) + d]$) the evaluation takes place:*

$$\rho'(\lambda, \varepsilon) = d^{-2} \exp(-\frac{4|\lambda|^{3/2}}{3\varepsilon}).$$

3) *If $\exp(-\frac{\operatorname{ctg}^3 \alpha}{3\varepsilon}) \leq d \leq \varepsilon^2$, then*

$$\int_{I_{d,\varepsilon}} d\rho(\lambda) = \frac{2 \operatorname{ctg} \alpha}{\sin^2 \alpha} + O(\varepsilon)$$

Corollary. *In space $C^*(I)$ the family of functions $\rho'(\lambda, \varepsilon)$ converges weakly to $\frac{2 \operatorname{ctg} \alpha}{\sin^2 \alpha} \delta(\lambda - \lambda_0)$ as $\varepsilon \rightarrow +0$.*

REFERENCES

1. Weyl H., *Über gewöhnliche Differentialgleichungen mit Singularitäten und die zugehörige Entwicklungen willkürlicher Funktionen*, Math. Ann. (1910), no. Bd. 68, 220-269.
2. Weyl H., *Über gewöhnliche lineare Differentialgleichungen mit singulären Stellen und ihre Eigenfunktionen*, Nachr. Acad. Wiss. Göttingen, Math. Phys. Klasse (1909), 37-64.
3. Weyl H., *Über gewöhnliche Differentialgleichungen mit singulären Stellen und ihre Eigenfunktionen*, Nachr. Acad. Wiss. Göttingen, Math. Phys. Klasse (1910), 442-467.
4. Levitan B.M., Sargsyan I.S., *Introduction to spectral theory* (1970), Moscow, Nauka. (Russian)
5. Levitan B.M., *Inverse problems of Sturm — Liouville* (1984), Moscow, Nauka. (Russian)
6. Marchenko V.A., *Sturm-Liouville operators and their applications* (1977), Kiev, Naukova Dumka. (Russian)
7. Titchmarsh E.C., *Some theorems on perturbation theory III*, Proc. Roy. Soc. **207** (1951, s.A), 321-328.

ALEXANDER S. PECHENTSOV AND ANTON YU. POPOV, MOSCOW STATE UNIVERSITY, RUSSIA.